

1.1 lakh women in India die every year due to causes related to pregnancy and child birth. Current strategies and programmes like Reproductive and Child Health Programme have therefore been addressed primarily to bringing about a reduction in the MMR and Infant and neonatal mortality.

Women above the age of 45 years form about 17% of the population. Evidence is also increasingly becoming available that hormonal and metabolic disorders and micro-nutrient deficiencies like Vitamin-D, cervical and breast cancers contribute to morbidity among women. Services for these ailments are being provided through the hospitals and other health care facilities. The Working Group on "Health Care for Women and Children" constituted by Planning Commission for Tenth Five Year Plan has also recommended adoption of a holistic life cycle approach towards women health, menopause, infertility, cervical and breast cancer etc. The National Population Policy-2000 also highlights the need for sensitizing, training and equipping rural need for sensitizing, training and equipping rural hospitals and health centres for providing geriatric care.

Survey under national programme for control of Blindness

499. SHRI RAMA MUNI REDDY SIRIGIREDDY: Will the Minister of HEALTH AND FAMILY WELFARE be pleased to state:

(a) whether the survey under the National Programme for Control of Blindness has been completed;

(b) if so, the details of the survey; and

(c) the details of the States from which data has been collected?

THE MINISTER OF STATE IN THE MINISTRY OF HEALTH AND FAMILY WELFARE (SHRI A. R&JA): (a) Yes, Sir.

(b) and (c) The survey has been undertaken in 15 randomly selected districts, one each from the States of Himachal Pradesh, Punjab, Uttar Pradesh, Bihar, West Bengal, Orissa, Madhya Pradesh, Rajasthan, Gujarat, Maharashtra, Chattisgarh, Andhra Pradesh, Karnataka, Tamil Nadu and Kerala. A total 72,008 persons aged 50 years and above were examined to find out prevalence and causes of blindness. The prevalence rate was 8.3% in population aged 50 years and above. Estimated prevalence of blindness in general population is 1.08%. Important causes of blindness were Cataract (63.5%) and Refractive Errors (27.6%).